

12015

DRAFT

Olivine Vitrophyre

191.2 grams




Figure 1: Photo of 12015 showing large vesicle (~3 cm dia.). Sample is 6 cm across.
NASA #S74-34503.

Introduction

12015 is an olivine vitrophyre generally similar to 12009. It is black because of its fine grained opaque matrix and also has a portion of a very large gas bubble (figure 1). The bulk composition of 12015 (and 12009) is thought to represent the original magma composition of Apollo 12 mare basalts.

Petrography

12015 is an olivine vitrophyre with skeletal and dendritic olivine and pyroxene phenocrysts (Baldrige et al. 1979). Microphenocrysts of chromite are also an early phase. These phenocrysts are set in a nearly opaque fine-grained matrix of dendritic pyroxene, plagioclase, filamental ilmenite, cristobalite, troilite, Fe-metal and glass.




Figure 2: Copies of photomicrographs of two thin sections of 12015 (from Baldrige et al. 1979).

Two thin sections of 12015 studied by Baldrige et al. (1979) had different textures (figure 2), both containing olivine phenocrysts set in fine-grained opaque matrix.

Mineralogy

Olivine: Olivine phenocrysts (Fo₇₆ to Fo₅₉) occur as equant to elongate grains <1 mm in size with slot-shaped inclusions of matrix. Rims of olivine contain chromite and metallic iron inclusions.

Pyroxene: Pyroxene compositions in 12015 were determined by Baldrige et al. (1979) (figure 3). The pyroxene compositions are very aluminous (> 9 wt. %)

Chromite: Chrome spinel is an early forming phase in 12015 (Baldrige et al. 1979).

Chemistry

The bulk chemical composition of 12015 appears to be similar to that of 12009 (table 1, figure 4 and 5). However, Baldrige et al. (1979) reported that the composition of “12015 lies on the extension of the fractionation trend defined by pigeonite basalts.” Baldrige et al. noted that removal of olivine and chromite from 12015 liquid composition could account for the composition of 12011 and 12043.

Walker et al. (1976), Rhodes et al. (1977) and Lindstrom and Haskin (1978) discuss the composition of 12015 (also 12009) as the starting magma composition of the Apollo 12 olivine basalt series. The olivine basalts correspond to a simple mixture of phenocryst olivine and 12015-like liquid.




Figure 3: Pyroxene and olivine composition of 12015 (adapted from Baldrige et al. 1979).

Radiogenic age dating

Nyquist et al. (1979) and Snyder et al. (1997) reported the isotopic composition of Sr and Nd, but it was not possible to determine an isochron age.

Other Studies

Bogard et al. (1971) reported the content and isotopic composition of rare gases in 12015.

Processing

12015 broke into three large pieces. There are 6 thin sections.

Mineralogical Mode for 12015

	Neal et al. 1994	Baldrige et al. 1979	
Olivine	62.3	10.3	20.4
Pyroxene		43	13.6
Plagioclase			
Ilmenite			
Chromite +Usp	3.2	0.3	0.2
mesostasis	33.7	46.3	65.7

List of Photo #s for 12015

S69-23391-393 B&W
 S69-62872-873
 S69-63342-350
 S69-64103
 S69-64128
 S70-24713-721
 S74-34502-503




Figure 4: The composition of 12015 compared with that of other lunar basalts.





Figure 5: Normalized rare-earth-element pattern for 12015 (from Nyquist 1979).



Table 1. Chemical composition of 12015.

reference weight	Neal 94 .548 g	Rhodes77	Nyquist79 41 mg	LSPET70	Baldridge79	Snyder97
SiO ₂ %		44.98 (c)		38	47	(d) 45
TiO ₂	2.9	(a) 2.86 (c)		3.2	2.9	(d) 2.86
Al ₂ O ₃	8.7	(a) 8.57 (c)		11	9.26	(d) 8.57
FeO	20.5	(a) 20.18 (c)		22	18.2	(d) 20.2
MnO	0.269	(a) 0.29 (c)		0.33	0.27	(d) 0.29
MgO	12.3	(a) 11.88 (c)		14	11.44	(d) 11.9
CaO	8.9	(a) 9.21 (c)		9.8	10.02	(d) 9.21
Na ₂ O	0.239	(a) 0.23 (a)		0.37	0.22	(d) 0.23
K ₂ O	0.06	(a) 0.06 (c)	0.054	(b) 0.062	0.05	(d) 0.06
P ₂ O ₅		0.06 (c)			0.06	(d) 0.06
S %		0.07 (c)			0.1	(d)
sum						
Sc ppm	48.4	(a) 46.1 (a)		44		47 (e)
V	186	(a)		95		
Cr	4250	(a) 4600 (a)		3900		2470 (e)
Co	47.8	(a) 51 (a)		47		51.9 (e)
Ni	62	(a) 50 (a)		70		73.5 (e)
Cu						12.4 (e)
Zn						12 (e)
Ga						3.63 (e)
Ge ppb						
As						
Se						
Rb			1.05	(b) 1		1.094 (e)
Sr	84	(a) 94 (c)	98.4	(b) 115		102.1 (e)
Y		35 (c)		46		34.5 (e)
Zr		110 (c)		160		127.6 (e)
Nb		6.6 (c)				8.01 (e)
Mo						
Ru						
Rh						
Pd ppb						
Ag ppb						218 (e)
Cd ppb						
In ppb						
Sn ppb						
Sb ppb						
Te ppb						
Cs ppm						0.078 (e)
Ba	65	(a) 94 (b)	60.1	(b) 44		67 (e)
La	6.2	(a)				6 (e)
Ce	16	(a) 16.3 (a)	16.1	(b)		16.7 (e)
Pr						2.7 (e)
Nd	14.4	(a)	12.2	(b)		16.1 (e)
Sm	4.3	(a) 4.31 (a)	4.14	(b)		4.77 (e)
Eu	0.98	(a) 0.81 (a)	0.869	(b)		1.07 (e)
Gd			5.6	(b)		6.1 (e)
Tb	1.16	(a) 1.05 (a)				1.1 (e)
Dy	7.4	(a)	6.7	(b)		6.98 (e)
Ho						1.5 (e)
Er			4.07	(b)		4.09 (e)
Tm						0.57 (e)
Yb	3.6	(a)	3.52	(b)		3.59 (e)
Lu	0.52	(a) 0.53 (a)	0.486	(b)		0.5 (e)
Hf	3.3	(a) 3.5 (a)				
Ta	0.38	(a)				0.432 (e)
W ppb						
Re ppb						
Os ppb						
Ir ppb						
Pt ppb						
Au ppb						
Th ppm	0.74	(a)				0.68 (e)
U ppm						0.33 (e)

technique (a) INAA, (b) IDMS, (c) XRF, (d) from mode, (e) ICP-MS

